

# The Gravid Surgical Patient

Chanel T. Tyler, MD  
 Assistant Professor  
 Obstetrics & Gynecology  
 Division of Maternal – Fetal Medicine  
 University of Wisconsin School of Medicine and Public Health


Bimbach, DJ Textbook of Obstetric Anesthesia, 2000

Parameter	Change	Amount %
Heart Rate	Increased	20-30
Stroke Volume	Increased	20-50
Cardiac Output	Increased	30-50
Contractility	Variable	+/-10
Central Venous Pressure	Unchanged	
Pulmonary Capillary Wedge Pressure	Unchanged	
Systemic Vascular Resistance	Decreased	20
Systemic Blood Pressure	Slight Decrease	Midtrimester 10-15 mm Hg, then rises
Pulmonary Vascular Resistance	Decreased	30
Pulmonary Artery Pressure	Slight Decrease	

## Objectives

- Understand Maternal Physiologic Adaptations to Pregnancy
- Understand Radiology Safety Concerns During Pregnancy
- Understand Anesthesia Safety Concerns
- Understand Preterm Labor in the Surgical Setting

## Cardiovascular Changes



- Plasma volume increases by 50%
- Red cell mass increases by 30%
- Dilutional anemia
- Uteroplacental shunt of 20-30%

## Maternal Physiologic Changes


## Cardiovascular Changes




- Systolic and diastolic BP nadir at 28 weeks
- SVR decreases during gestation
- Supine hypotension

## Supine Hypotension

Figure 1: Supine Hypotensive Syndrome




## Changes in Fibrinogen in Pregnancy



## Coagulation Changes in Pregnancy

- Normal pregnancy is associated with major changes in all aspects of hemostasis
- Various aspects of the DIC diagnostic panel are affected
- There are changes from trimester to trimester along with during labor, delivery and the postpartum period


## Changes in Antithrombin in Pregnancy



## Hematopoietic System

- ↑ Plasma volume
- Increased RBC production 15-20%
  - Dilutional anemia
- Leukocytosis (6-16)
- ↑ fibrinogen, V, VII, VIII, IX, X, XII
- ↓ XI, XIII

## Changes in D-dimer in Pregnancy




## Respiratory Changes

- Compensated respiratory alkalosis
- Lowered  $P_a\text{CO}_2$
- Lowered bicarbonate (18-22 mEq/L)
- Minute ventilation increases by 40%
- Decreased functional residual volume
- Elevated diaphragm by 4 cm

## Alterations in Anatomy

- 1<sup>st</sup>-trimester uterus is thick-walled and intra-pelvic
- Out of pelvis > 12 wks
- 2<sup>nd</sup>-trimester uterus contains large amount of amniotic fluid
- 3<sup>rd</sup>-trimester uterus is thin-walled, large; fetal head engaging pelvis
- At 36 weeks, uterus reaches costal margin



## Blood Gas Values

pH	7.40-7.45
pO <sub>2</sub>	104-108 mm Hg
pCO <sub>2</sub>	27-32 mm Hg
HCO <sub>3</sub>	18-22 mEq/L



## Kidneys and Urinary Tract



- Renal enlargement, dilation and position change
- GFR increases by 50%
- Impaired tubular glucose resorption
- Bladder is displaced upward and 'flattened'

## GI Changes

- Reduced motility (controversial)
- Large and small bowel move upward and laterally
- Decreased gallbladder emptying





## Diagnostic Studies

### ACOG Committee Opinion Number 299, September 2004

- Ultrasonography and MRI are not associated with known adverse fetal effects.
- Consultation with an expert in dosimetry calculation may be helpful in calculating estimated fetal dose when multiple diagnostic X-rays are performed on a pregnant patient.

(Replaces No. 158, September 1995) (Reaffirmed 2009) Committee on Obstetric Practice

### ACOG Committee Opinion Number 299, September 2004

- Women should be counseled that X-ray exposure from a single diagnostic procedure does not result in harmful fetal effects. Specifically, exposure to less than 5 rad has not been associated with an increase in fetal anomalies or pregnancy loss.

(Replaces No. 158, September 1995) (Reaffirmed 2009) Committee on Obstetric Practice

### ACOG Committee Opinion Number 299, September 2004

- The use of radioactive isotopes of iodine is contraindicated for therapeutic use during pregnancy.
- Radiopaque and paramagnetic contrast agents are unlikely to cause harm and may be of diagnostic benefit, but these agents should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

(Replaces No. 158, September 1995) (Reaffirmed 2009) Committee on Obstetric Practice

### ACOG Committee Opinion Number 299, September 2004

- Concern about possible effects of high-dose ionizing radiation exposure should not prevent medically indicated diagnostic X-ray procedures from being performed on a pregnant woman.
- During pregnancy, other imaging procedures not associated with ionizing radiation (i.e., ultrasonography, MRI) should be considered instead of X-rays, when appropriate.

(Replaces No. 158, September 1995) (Reaffirmed 2009) Committee on Obstetric Practice

## Plain Films

- **< 5-10 rads**
  - No risk on congenital malformation, abortions or intra-uterine growth ret.
  - Smaller risk of increase in childhood cancer
- **Radiation doses > 10 rads**
  - 6 % chance of severe mental impairment
  - < 3 % chance childhood cancer

Rosenstein, HEW publication (FDA) 89-8031, Rockville, MD, US Dept. Of Health And Human Services, Centre For Devices And Radiologic Health, 1988.

Radiographic Examination	Dose to Ovary / Uterus-mrad
<b>Low-Dose Group:</b>	
Head, C-Spine, Thoracic Spine Chest, Extremities	<1
<b>High-Dose Group:</b>	
Lumbar Spine	204 – 1260
Pelvic	190 – 357
Hip	124 – 450
Intravenous Pyelogram	503 – 880
Urethrocytogram	1500
KUB	200 – 503

Rosenstein M-Handbook of selected organ doses for projections common in diagnostic radiology, HEW publication(FDA) 89-8031, Rockville MD, US Dept. Of Health And Human Services, Centre For Devices And Radiologic Health, 1988.

Radiographic Examination	Dose (mrad)
<b>Computed Tomography</b>	
Head (1 cm slice)	< 50
Chest (1 cm slice)	< 1000
Upper Abdomen (20 slices 2.5 cm above uterus)	< 3000
Lower Abdomen (10 1 cm slices over the uterus/ fetus)	3000 – 9000
<b>Angiography</b>	
Cerebral	< 100
Cardiac Catheterization	< 500
Aortography	< 100

- ## Ultrasound
- Best modality to assess both fetus and mother
  - Not sensitive:
    - Colonic lesions
    - Biliary tree lesions
    - Sub-placental hematoma
  - Safe procedure

**Bottom Line:**

**If the study is needed to make the diagnosis  
DON'T COMPROMISE!**

- ## CT SCAN
- Complementary to U/S & DPL
  - Penetrating wounds of flank & back
  - Can miss diaphragmatic and bowel injuries
  - portability
  - Spiral CT reduces radiation exposure by 14-30 %

**REMEMBER:**

**A POORLY TREATED MOTHER = A POORLY TREATED BABY!!!!**

## Anesthesia Concerns

- Progesterone relaxes the smooth muscle
  - Impairs esophageal and intestinal motility during pregnancy
  - Delayed gastric emptying is controversial
    - However, the risk of pulmonary aspiration of gastric contents remains real
      - Endotracheal intubation reduces this risk

## Anesthesia Concerns

- Capillary engorgement of the mucosa and edema of the oropharynx, larynx, trachea may lead to difficult intubation
- Friable mucosa
  - Intubation may cause bleeding making visualization while securing airway difficult

## Anesthesia Concerns

Maternal accommodation of the increased oxygen demands and requirement for carbon dioxide elimination:

- Decrease in functional residual capacity
- Increase minute ventilation as a result of increased tidal volume
- Unchanged respiratory rate
- Unchanged closing capacity
  - Therefore, decreased FRC/CC ratio
  - Faster airway closure with reduced lung volume

## Anesthesia Concerns

- Uterine Blood Flow
  - Lacks autoregulation as vessels are maximally dilated
  - Uterine artery flow is dependent on maternal blood pressure and cardiac output

**Bottom Line: ALTER MATERNAL BLOOD PRESSURE, ALTER FETAL BLOOD SUPPLY**

## Anesthesia Concerns

**Bottom Line: PREGNANT WOMEN WILL DESATURATE FASTER THAN NON-PREGNANT WOMEN**

**Solution: ADMINISTRATION OF 100% O<sub>2</sub> FOR 3-5 MINUTES PRIOR TO INDUCTION OF ANESTHESIA (capacity breaths in and emergency situation sufficient)**

## Anesthesia Concerns

- Pregnant women demonstrate increased sensitivity to both regional and general anesthesia
- Extent of drug transfer is dependent on numerous factors
  - Molecular weight
  - Degree of lipid solubility
  - Maternal drug concentration
  - Maternal and Fetal pH

- Lower fetal pH favors ionization of basic local anesthetics (i.e., lidocaine) and may explain the accumulation of drugs in a compromised fetus

## Differential Diagnosis

- Appendicitis
- Ovarian Torsion
- Cholecystitis
- Cholelithiasis
- Pancreatitis
- Degenerating Fibroid



## Intra- OP

- Uneventful LMA placement
- Laparoscopic Appendectomy
- Transferred to PACU
  - No contractions
  - FHR 155 by Doptone

## Case

- 24 yo G1P0 @ 18 wks uncomplicated
- c/o generalized abdominal pain 3-hr duration, nausea vomiting, afebrile, no vaginal bleeding
- UA neg, WBC 16,000
- Fetal ultrasound without defects
- FHR 160 by Doptone

## 1-Hour Postop

- Reports shortness of breath and chest pain
  - HR 105
  - Pox 85%
  - Get ABG
    - 7.3/80/40

- EKG
  - Right axis deviation and right bundle branch block
  - What's normal
- CXR
  - Apparent cardiomegaly with increased vascular markings

## What's Normal in Pregnancy?

	Pregnant	Pre-Pregnant
pH	7.4-7.45	7.35-7.45
PO <sub>2</sub> (mmHg)	104-108	80-100
PCO <sub>2</sub> (mmHg)	27-32	35-42
HCO <sub>3</sub> (mEq/L)	22	26

- Compensated respiratory alkalosis
  - PaCO<sub>2</sub> of 40mmHg is very abnormal in pregnancy
  - Fetus relies on high maternal PaO<sub>2</sub>

## What Concerns You Most?

- Tachycardia
- Pulse OX
- ABG
- CXR
- EKG
- Echo

## Normal CXR in Pregnancy

- Apparent Cardiomegaly
- Enlarged left atrium (lateral views)
- Increased vascular markings
- Straightening of left-sided heart border
- Postpartum pleural effusion

Modified from Gei AF, Hawkins GDV. Cardiac disease and pregnancy. Obstet Gynecol Clin North Am 28: 465-512, 2001

## What Concerns the Obstetrician Most?



ABG

## Normal Electrocardiography

- Right axis deviation
- Right bundle branch block
- ST segment depression of 1 mm on left precordial leads
- Q waves in lead III
- T-wave inversion in leads III, V<sub>2</sub>, V<sub>3</sub>
- Small decrease in PR and QT interval (heart-rate dependent)
- Rotation +/- 25 degrees (QRS axis)

Modified from Gei AF, Hawkins GDV. Cardiac disease and pregnancy. Obstet Gynecol Clin North Am 28: 465-512, 2001

## Normal Echocardiography

- Trivial tricuspid regurgitation (up to 43-93% at term)
- Pulmonary regurgitation (up to 94% at term)
- Increased atrial size 12-14%
- Increased left ventricle end-diastolic dimensions by 6-10%
- Inconsistent increase in left ventricle thickness
- Mitral regurgitation (28% at term)
- Pericardial effusion (40% postpartum)

Remember:

**What is Best  
for the Mother is  
always Best for  
the Fetus!**

When to Intervene and  
Consult

**EARLY!**

Remember:

**YOU CANNOT HAVE A  
HEALTHY BABY  
WITHOUT A HEALTHY  
MOTHER !**

Remember:

**Delivery is NOT  
Always in the  
Mothers Best  
Interest!**




## Acute Hemorrhage

	Class 1	Class 2	Class 3	Class 4
Blood loss % blood volume	15%	15-30%	30-40%	>40%
Pulse rate	<100	>100	>120	>140
Pulse pressure	Normal	Decreased	Decreased	Decreased
Blood Pressure	Normal or increased	Decreased	Decreased	Decreased

## Fetal Physiology

- Gestational age dependent
- Severity and type of trauma
- Normal fetal physiology




## Teratology Concerns

- Radiation exposure
- Placental metabolism
- Category X drugs
- Variation according to gestational age

## Modalities for Evaluating Trauma

- Plain Films – X-rays
- Ultrasound
- CT & MRI
- Cardiocotographic Monitoring
- DPL
- Laparotomy



## Diagnostic Peritoneal Lavage

- CT & U/S are better in stable patient
- Can be performed in any trimester
- Gravid uterus does not reduce the accuracy Limited in detecting bowel perforation and does not assess retroperitoneal hemorrhage or intra-uterine pathology

Rotherberg DA, et al: Diagnostic peritoneal lavage for blunt trauma in pregnant women. Am J Obstet Gynecol 129:479-481, 1977.

Epstein TJ, et al: Evaluation of blunt abdominal trauma occurring during pregnancy. J Trauma 29:1628-1632, 1989.